

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 JAN 2009		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Substrate utilization in the critically ill				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Wolf S. E., Shields B. A., Wade C. E.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Journal of Parenteral and Enteral Nutrition

<http://pen.sagepub.com/>

Letter to the Editor

Steven E. Wolf, Beth A. Shields and Charles E. Wade
JPEN J Parenter Enteral Nutr 2009 33: 111
DOI: 10.1177/0148607108319798

The online version of this article can be found at:
<http://pen.sagepub.com/content/33/1/111>

Published by:



<http://www.sagepublications.com>

On behalf of:



American Society for Parenteral
and Enteral Nutrition

[The American Society for Parenteral & Enteral Nutrition](#)

Additional services and information for *Journal of Parenteral and Enteral Nutrition* can be found at:

Email Alerts: <http://pen.sagepub.com/cgi/alerts>

Subscriptions: <http://pen.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [Version of Record](#) - Jan 22, 2009

[What is This?](#)

Dear Editor,

We read the recent paper Effect of Calorically Dense Enteral Nutrition Formulas on Outcome in Critically Ill Trauma and Surgical Patients¹ with some interest, as this has been an investigative interest of our group. The findings are very intriguing and fit our bias regarding substrate utilization in the critically ill. One of the key and perhaps most compelling differences between normocaloric and calorically dense formulae is simply the fat content; the only method to increase caloric density is to increase the lipid component in relation to carbohydrate and protein. In this study, close to double the amount of fat was delivered to the calorically dense group. Substrate delivery and utilization during critical illness, then, comes to the forefront in deciphering the results of this study. Should fat be delivered as the primary substrate, and is this beneficial or harmful? It was shown previously that glucose/carbohydrate appears to be the preferential substrate in critical illness related to poor utilization of fat with impaired oxidation and inefficient transport between pools,^{2,3} which would argue against the use of fat in this population as a testable hypothesis.

We showed several years ago that low fat feeding is associated with better muscle protein accretion in a severely burned surgical population compared with an isonitrogenous isocaloric high-fat feeding, which was associated with increased endogenous insulin release.⁴ Garrel and others showed that low-fat enteral feeding was associated with improved clinical outcomes in a similar population.⁵ This current study now concludes that a lower fat formula is also associated with better clinical outcomes in a critically ill nonburned surgical population, providing further rationale that lower fat feeding should be considered over high-fat feeding in any critically ill surgical patient.

That being stated, this study does have some problems in design from variables that were not measured. The difference in choice of feeding regimens was likely associated with provider preference; therefore, the measured effects may be more associated with the provider than the formula. For instance, what about ventilator-weaning protocols, were they similar between providers? In addition, were these data gathered before the wide acceptance of glucose control with aggressive insulin therapy? Is it possible there is a role of insulin dosing in the normocaloric group?

In conclusion, we are happy to see the results of this study, which we propose is related to bulk substrate delivery and utilization, and not some micronutrient effect. The data presented are enticing, but by no means definitive. We wholeheartedly agree with the authors that a prospective trial is needed to assess bulk substrate utilization with a directed examination of delivery of differing percentages of the primary energy substrates of carbohydrate and fat upon overall clinical effects. Specific effects on muscle, liver, fat, and the immune system should also be examined.

Yours sincerely,

Steven E. Wolf, MD
Department of Surgery
University of Texas Health Science Center—San Antonio
Clinical Research, Burn Center
United States Army Institute of Surgical Research

Beth A. Shields, RD
United States Army Institute of Surgical Research
Brooke Army Medical Center

Charles E. Wade, PhD
United States Army Institute of Surgical Research

References

1. Bryk J, Zenati M, Forsythe R, Peitzman A, Ochoa JB. Effect of calorically dense enteral nutrition formulas on outcome in critically ill trauma and surgical patients. *JPEN J Parenter Enteral Nutr.* 2008;32:6-11.
2. Cree MG, Aarsland A, Herndon DN, Wolfe RR. Role of fat metabolism in burn trauma-induced skeletal muscle insulin resistance. *Crit Care Med.* 2007;35(9 Suppl):S476-S483.
3. Aarsland A, Chinkes DL, Sakurai Y, Nguyen TT, Herndon DN, Wolfe RR. Insulin therapy in burn patients does not contribute to hepatic triglyceride production. *J Clin Invest.* 1998;101:2233-2239.
4. Hart DW, Wolf SE, Zhang XJ, et al. Efficacy of a high-carbohydrate diet in catabolic illness. *Crit Care Med.* 2001;29:1318-1324.
5. Garrel DR, Razi M, Lariviere F, et al. Improved clinical status and length of care with low-fat nutrition support in burn patients. *JPEN J Parenter Enteral Nutr.* 1995;19:482-491.